


2023

4th Semester Examination  
PHYSICS (Honours)

Paper : C 8-T

[Mathematical Physics - III]

[CBCS]



Full Marks : 40

Time : Two Hours

*The figures in the margin indicate full marks.  
Candidates are required to give their answers  
in their own words as far as practicable.*

Illustrate the answer wherever necessary.  
Symbols have their usual meaning.

1. Answer any **five** from the following :  $2 \times 5 = 10$

(i) Find the square root of the following  $3 + 4i$ .

(ii) Prove that  $(AB)^{-1} = B^{-1}A^{-1}$ .

(iii) In which domain(s) of the complex plane  
 $f(z) = |x| - i|y|$  is an analytic function?

(iv) Identify the zeroes, poles and essential singularities  
of  $e^{\frac{1}{z}}$ .

P.T.O.

( 2 )

(v) Simplify the expression  $z = i^{-2i}$ .

(vi) What is the Fourier transform of Dirac delta  $\delta(x-x_0)$ ?

(vii) Write convolution theorem involving Fourier transform.

(viii) Prove eigenvalues of a hermitian matrix are real.

2. Answer any *four* from the following :  $5 \times 4 = 20$

(i) Prove that  $\int_0^{2\pi} \frac{d\theta}{2 + \cos\theta} = \frac{2\pi}{\sqrt{3}}$ .

(ii) Find the Fourier transform of the normalised Gaussian distribution

$$f(t) = \frac{1}{\tau\sqrt{2\pi}} e^{-\frac{t^2}{2\tau^2}} \quad -\infty \leq t \leq \infty$$

(iii) Find a function  $f(z)$ , analytic in a suitable part of the Argand diagram, for which

$$\operatorname{Re} f = \frac{\sin x}{\cosh 2y - \cos 2x}$$

Where are the singularities of  $f(z)$ ?

(iv) Determine the types of singularities (if any) possessed by the following functions at  $z = 0$  and  $z = \infty$  :

( 3 )

(a)  $(z-2)^{-1}$ ,

(b)  $(1+z^3)/z^2$ ,

(c)  $\sinh(1/z)$ ,

(d)  $e^z/z^3$ .

(v) What is Cauchy Riemann condition? Apply on the function  $f(z) = |z|^2$  and comment on its analyticity.

(vi) Using Cayley Hamilton Theorem find the inverse matrix

$$\begin{vmatrix} \cos A & \sin A \\ -\sin A & \cos A \end{vmatrix}$$

3. Answer any *one* from the following :  $10 \times 1 = 10$

(i) (a) Find the Fourier transform of the given function

$$f(x) = 1 \text{ for } |x| < a \\ 0 \text{ for } |x| > a.$$

(b) Using contour integration to evaluate the real integral

$$\int_0^{\infty} \frac{1}{1+x^2} dx = \frac{\pi}{2}.$$

P.T.O.




Fig.1

Find the Fourier transform of the function in figure 1 representing two wide slits by considering the Fourier transforms of

- (A) two  $\delta$ -functions, at  $x = \pm a$ ,
- (B) a rectangular function of height 1 and width  $2b$  centred on  $x = 0$ .  $2+2=4$
- (b) Find the Fourier transform of the function  $f(t) = \exp(-|t|)$ .

(A) By applying Fourier's inversion theorem prove that

$$\frac{\pi}{2} \exp(-|t|) = \int_0^{\infty} \frac{\cos \omega t}{1 + \omega^2}.$$

(B) By making the substitution  $\omega = \tan \theta$ , demonstrate the validity of Parseval's theorem for this function.  $3+3=6$